ELSEVIER

Journal of Organometallic Chemistry 506 (1996) 363–364

Journal of Organometallic Chemistry

Author Index of Volume 506

Adams, C.J., 191
Akhnoukh, T., 113
Akiba, K.-Y., 259
Al-Jibori, S.A., 119
Alvarez-Larena, A., 121
Alvarez Boo, P., 253

Baratta, W., 357
Bardin, V.V., 155
Bharadwaj, S.K., 181
Blenkiron, P., 229
Bosch, W.H., 273
Breckenridge, S.M., 229
Bruce, M.I., 191

Cai, R.-F., 25
Caiazzo, A., 337
Carty, A.J., 229
Caruso, F., 67
Casas, J.S., 253
Castiñeiras, A., 253
Cerrada, E., 203
Chadha, R.K., 181
Chang, Y.-T., 101
Cheung, K.-K., 77
Cloke, F.G.N., 343
Consiglio, G., 177
Couce, M.D., 253

Dash, A.K., 307
Deacon, G.B., 1
Duncalf, D.J., 347

Englert, U., 273

Ferguson, G., 129
Fernández, A., 165
Fernández, J.J., 165
Frank, W., 293
Freijanes, E., 253
Frohn, H.J., 155
Fu, P.-F., 49

Gayoso, M., 165
Geldbach, T.J., 343
Germain, G., 121
Giesen, M., 155
Gimeno, M.C., 203
Giomini, M., 67
Giorgi, G., 61
Girdwood, S.E., 321
Giuliani, A.M., 67
Glidewell, C., 129
Grady, A.S., 327
Gräfe, A., 31
Green, M.L.H., 221
Guillon, C., 211

Harris, P.W.R., 339
Hassler, K., 93
Haupt, H.-J., 41
Hayashi, T., 85
Herrmann, W.A., 351, 357
Hiller, J., 109, 241
Hitchcock, P.B., 343, 347
Holm, T., 37
Hong, F.-E., 101
Hossain, Md.M., 307, 313
Huang, X.-Y., 25
Huang, Z.-E., 25
Hughes, A.K., 221

Ito, Y., 85

Jha, N.K., 19
Jones, P.G., 203

Kaur, S., 181
Khan, M.A., 49
Klose, A., 155
Korrell, G., 293
Kruck, T., 31
Kunkely, H., 175

Laguna, A., 203
Laguna, M., 203
Lawless, G.A., 347
Lazzaroni, R., 337
Lee, S.-M., 77
Lewin, A., 155
Lin, C.-C., 101
Linney, R.E., 327
Liou, M.-W., 101
Lo, S.-C., 101
López-Solera, I., 149
López Torres, M., 165
Louattani, E., 121
Love, J.B., 343

Mach, K., 109, 241
Macleod, I.T., 301
Mahmood, Z., 327
Markwell, R.D., 327
Masaguer, J.R., 149
Mathur, P., 307, 313
Matković-Čalogović, D., 265
Meider, H., 265
Monge, A., 149
Morawietz, M.J.A., 351
Muir, K.W., 321
Müller, J., 113

Narula, S.P., 181
Navarro-Ranninger, C., 149
Nicholas, K.M., 49

Ohashi, N., 259
Opromolla, G., 61, 129
Orera, V., 203
Ortigueira, J.M., 165

Pellinghelli, M.A., 229
Pereira, T., 165
Pétillon, F.Y., 321
Pfister, B., 273
Piniella, J.F., 121
Planinić, P., 265
Poder-Guillou, S., 321
Polášek, M., 241
Pöschl, U., 93
Priermeier, T., 351
Pritzkow, H., 287
Puranik, V.G., 313

Reiß, G.J., 293
Rivarola, E., 67
Russell, D.K., 327
Russo, U., 253

Sakaguchi, A., 259
Salzer, A., 273
Sánchez González, A., 253
Sappa, E., 229
Sasso, G., 61
Satyanarayana, C.V.V., 307, 313
Schmidt, Th., 139
Schniedermeier, J., 41
Schollhammer, P., 321
Sedmera, P., 241
Settambolo, R., 337
Shankar, R., 181
Sharma, P., 19
Shen, Q., 1

Siegl, H., 93
Skelton, B.W., 191
Sonn, I., 113
Sordo, J., 253
Sperrle, M., 177
Stauber, R., 273
Suades, J., 121

Talarmin, J., 321
Tamao, K., 85
Taylor, N.J., 229
Thewalt, U., 109, 241, 331
Tiekink, E.R.T., 301
Tiripicchio, A., 229
Togni, A., 61
Troyanov, S.I., 241

Umbarkar, S., 313

Varga, V., 109, 241
Verghese, B., 307
Vierling, P., 211
Vila, J.M., 165
Vogler, A., 175

Wadepohl, H., 287
White, A.H., 191
Wöhrle, T., 331
Wolf, A., 287
Wong, W.-T., 77
Woodgate, P.D., 339
Wu, Z.-Z., 25

Xu, Z., 25

Yamamoto, Y., 259
You, X.-Z., 25
Young, C.G., 301

Zakaria, C.M., 129
Zamora, F., 149
Zanello, P., 61, 129
Zhou, X.-G., 25

ELSEVIER

Journal of Organometallic Chemistry 506 (1996) 365–372

Journal of Organometallic Chemistry

Subject Index of Volume 506

(Trimethylsilyl)acetylene complexes
Titanocene-bis(trimethylsilyl)acetylene complexes: effects of methyl substituents at the cyclopentadienyl ligands on the structure of thermolytic products (V. Varga, K. Mach, M. Polášek, P. Sedmera, J. Hiller, U. Thewalt and S.I. Troyanov), 241

μ-η^1:η^2-Acetylide
Comparative chemistry of μ-η^1 : η^2-allenyl and μ-η^1 : η^2-acetylide complexes: η^5-cyclopentadienyl complexes via alkyne-allenyl coupling in the reactions of PhC≡CPh with $[Ru_2(CO)_6\{\mu$-η^1 : $\eta^2_{3,4}$-C(Ph)=C=$CH_2\}(\mu$-$PPh_2)]$ and $[Ru_2(CO)_6\{\mu$-η^1 : η^2-C≡CPh$\}(\mu$-$PPh_2)]$. X-Ray structures of $[\{Ru_2(CO)_4\{\mu$-η^2(C,O),η^5-$C_5MePh_2(C_6H_4O)\}(\mu$-$PPh_2)$$_2]$ and $[Ru_2(CO)_5$-$\{\mu$-η^2(C,O),η^5-$C_5HPh_2(C_6H_4O)\}(\mu$-$PPh_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229

Alkyl
Thermolysis of $(MeCp)_2Nb(\eta^2$-$CO_2)R$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49

Alkyne
CpCo-moderated stereospecific dimerization of $(Me_3SiC_2)_2S$ to give [*trans*-{$(Me_3Si)_2(Me_3SiC_2S)_2C_4$}CoCp]: synthesis and structure of [*trans*-{$(Me_3Si)_2[(Co_2(CO)_6(\mu$-$\eta^2$: η^2-$Me_3SiC_2S)]_2C_4\}$CoCp] (H. Wadepohl, A. Wolf and H. Pritzkow), 287

Alkyne–allenyl coupling
Comparative chemistry of μ-η^1 : η^2-allenyl and μ-η^1 : η^2-acetylide complexes: η^5-cyclopentadienyl complexes via alkyne-allenyl coupling in the reactions of PhC≡CPh with $[Ru_2(CO)_6\{\mu$-η^1 : $\eta^2_{3,4}$-C(Ph)=C=$CH_2\}(\mu$-$PPh_2)]$ and $[Ru_2(CO)_6\{\mu$-η^1 : η^2-C≡CPh$\}(\mu$-$PPh_2)]$. X-Ray structures of $[\{Ru_2(CO)_4\{\mu$-η^2(C,O),η^5-$C_5MePh_2(C_6H_4O)\}(\mu$-$PPh_2)$$_2]$ and $[Ru_2(CO)_5$-$\{\mu$-η^2(C,O),η^5-$C_5HPh_2(C_6H_4O)\}(\mu$-$PPh_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229

μ-η^1:η^2-Allenyl
Comparative chemistry of μ-η^1 : η^2-allenyl and μ-η^1 : η^2-acetylide complexes: η^5-cyclopentadienyl complexes via alkyne-allenyl coupling in the reactions of PhC≡CPh with $[Ru_2(CO)_6\{\mu$-η^1 : $\eta^2_{3,4}$-C(Ph)=C=$CH_2\}(\mu$-$PPh_2)]$ and $[Ru_2(CO)_6\{\mu$-η^1 : η^2-C≡CPh$\}(\mu$-$PPh_2)]$. X-Ray structures of $[\{Ru_2(CO)_4\{\mu$-η^2(C,O),η^5-$C_5MePh_2(C_6H_4O)\}(\mu$-$PPh_2)$$_2]$ and $[Ru_2(CO)_5$-$\{\mu$-η^2(C,O),η^5-$C_5HPh_2(C_6H_4O)\}(\mu$-$PPh_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229

Allyl halide
Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[CpMo(CO)_3H]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(C_5H_5)Mo(CO)(\mu$-$SMe)_3Mo(CO)_2(\mu$-$SMe)Mo(CO)_2(C_5H_5)]$ (S. Poder-Guillou, P. Schollhammer, F.Y. Pétillon, J. Talarmin, S.E. Girdwood and K.W. Muir), 321

Amide
First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357

Amino-functionalized cyclopentadienyl
First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357

Anion
Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin–carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259

Ansa-metallocenes
Unsymmetrical *ansa*-metallocenes of zirconium and hafnium (W.A. Herrmann, M.J.A. Morawietz and T. Priermeier), 351

Association
Association and exchange in dialkylgallanes in toluene solution (D.K. Russell, A.S. Grady, R.E. Linney, Z. Mahmood and R.D. Markwell), 327

Benzofuranylferrocene
Synthesis and characterization of new perfluoroalkylated side-chain ferrocenes and ferricinium salts (C. Guillon and P. Vierling), 211

Bidentate Lewis acid
Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85

Bidentate ligand
Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin–carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259

Bipyrimidine
Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2′-bipyrimidine and further NMR studies of diorganotin-pyrazine and −2,2′-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

Bis(arene) complexes
Schwermetall-π-Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293

Bridging thiolate
Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[CpMo(CO)_3H]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(C_5H_5)Mo(CO)(\mu$-$SMe)_3Mo(CO)_2(\mu$-$SMe)Mo(CO)_2(C_5H_5)]$ (S. Poder-Guillou, P. Schollhammer, F.Y. Pétillon, J. Talarmin, S.E. Girdwood and K.W. Muir), 321

Butadiene
A chalcogen-stabilized cluster containing a s-*trans*-1,3-diene ligand: synthesis and structural characterization of $\{(CO)_6Fe_2Se_2\}_2(\mu$-s-trans-$C_4H_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307

Carbon dioxide
Thermolysis of $(MeCp)_2Nb(\eta^2$-$CO_2)R$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49

Carbonyl
A chalcogen-stabilized cluster containing a s-*trans*-1,3-diene ligand: synthesis and structural characterization of $\{(CO)_6Fe_2Se_2\}_2(\mu$-s-trans-$C_4H_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307
Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyldiphenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of $[Cr(CO)_4]_2$TPTA and $[Cr(CO)_5]_4$TPTA (H. Meider, D. Matković-Čalogović and P. Planinić), 265
Propargyl alcohol derivatives of $Fe_2(CO)_6(\mu$-EE′) (E,E′ = S, Se or Te): synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu$-EC(H)=C(CH_2OH)E′} (E = Se or Te; E′ = S, Se or Te); Crystal structure of $(CO)_5(PPh_3)Fe_2\{\mu$-SeC(CH_2OH)=C(H)Se} (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313
Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121
Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortigueira), 165

Carbonyls
Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[HOs_3(CO)_{10}(O_2CC_5H_4FeC_5H_5)]$ and $[\{Os_3H(CO)_{10}\}_2(O_2CC_5H_4FeC_5H_4CO_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77

Carboxylate
Crystal and molecular structures of 1,2;3,4-Di-μ-*o*-methyl/*o*-chlorobenzoato-*OO*′-1,3-bis(o-methyl/o-chlorobenzoato-*O*)-1,2,4;2,3,4-di-μ_3-oxotetrakis[di-*n*-butyltin(IV)] and molecular structure of 1,2;3,4-Di-μ-*o*-hydroxybenzoato-*OO*′-1,3-bis(o-hydroxybenzoato-*O*)-1,2,4;2,3,4-di-μ_3-oxotetrakis[di-*n*-butyltin(IV)] compounds (S.P. Narula, S. Kaur, R. Shankar, S.K. Bharadwaj and R.K. Chadha), 181

Catalyses
Novel zirconium complexes derived from C_2-symmetric diamide ligands; the X-ray crystal structure of $[Zr(\eta^1$-$CH_2Ph)(\eta^2$-$CH_2Ph)\{(C_6H_3)_2$-2,2′-$(NCH_2C_6H_4Ph$-4$)_2$-6,6′-$Me_2\}]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343

Catalysis
High α-regioselectivity in the rhodium-catalyzed hydroformylation of vinylpyrroles (R. Settambolo, A. Caiazzo and R. Lazzaroni), 337
Ruthenium-catalysed *ortho* alkylation of hydroxyacetophenones; the functionalisation of ring C aromatic diterpenoids (P.W.R. Harris and P.D. Woodgate), 339

Catalysts
New rhodium(I)-π-chelate complexes with coordinated amidine bases (dbu, dbn) and their catalytic properties to polymerize phenylacetylene (J. Schniedermeier and H.-J. Haupt), 41

Charge–transfer salts
Charge-transfer salts with mononuclear and dinuclear ylide gold(I) complexes: X-ray structure of $[Au(CH_2PPh_3)_2](TCNQ)$ (TCNQ = 7,7′,8,8′-tetracyanoquinodimethane) (E. Cerrada, M.C. Gimeno, A. Laguna, M. Laguna, V. Orera and P.G. Jones), 203

Chromium
Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyldiphenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of $[Cr(CO)_4]_2$TPTA and $[Cr(CO)_5]_4$TPTA (H. Meider, D. Matković-Čalogović and P. Planinić), 265
Synthesis and characterization of tricarbonyl[η^6-(phenylethynyl)-arene]chromium complexes: crystal structure of tricarbonyl[η^6-(3-phenylethynyl)anisole]chromium (F.-E. Hong, S.-C. Lo, M.-W. Liou, Y.-T. Chang and C.-C. Lin), 101

Cluster
A chalcogen-stabilized cluster containing a s-*trans*-1,3-diene ligand: synthesis and structural characterization of $\{(CO)_6Fe_2Se_2\}_2(\mu$-s-trans-$C_4H_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307
Cluster chemistry XCVIII. Formation of an Ru_5–dicarbon complex: X-ray structures of $Ru_5(\mu_5$-$C_2)(\mu$-$PPh_2)_2(\mu$-$Cl)_2(CO)_{11}$ and $Ru_5(\mu$-$H)(\mu_5$-$CC(PPh_2))(\mu$-$Cl)(\mu$-$PPh_2)(CO)_{12}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 191
Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[HOs_3(CO)_{10}(O_2CC_5H_4FeC_5H_5)]$ and $[\{Os_3H(CO)_{10}\}_2(O_2CC_5H_4FeC_5H_4CO_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77

^{13}C NMR spectra
Tetrachloropyridylstibines (P. Sharma and N.K. Jha), 19

Cobalt
CpCo-moderated stereospecific dimerization of $(Me_3SiC_2)_2S$ to give $[trans$-$\{(Me_3Si)_2(Me_3SiC_2S)_2C_4\}CoCp]$: synthesis and structure of $[trans$-$\{(Me_3Si)_2[(Co_2(CO)_6)(\mu$-$\eta^2:\eta^2$-$Me_3SiC_2S)]_2C_4\}CoCp]$ (H. Wadepohl, A. Wolf and H. Pritzkow), 287
Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121
Reaktionen von Nitrosylkomplexen. XII. Umsetzung von $[\{C_5Me_5$-$Fe(\mu$-$NO)\}_2]$ mit $[C_5Me_5Co(C_2H_4)_2]$: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnoukh), 113

Coupling reaction
CpCo-moderated stereospecific dimerization of $(Me_3SiC_2)_2S$ to give $[trans$-$\{(Me_3Si)_2(Me_3SiC_2S)_2C_4\}CoCp]$: synthesis and structure of $[trans$-$\{(Me_3Si)_2[(Co_2(CO)_6)(\mu$-$\eta^2:\eta^2$-$Me_3SiC_2S)]_2C_4\}CoCp]$ (H. Wadepohl, A. Wolf and H. Pritzkow), 287

Crystal structure
A chalcogen-stabilized cluster containing a s-*trans*-1,3-diene ligand: synthesis and structural characterization of $\{(CO)_6Fe_2Se_2\}_2(\mu$-s-trans-$C_4H_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307
Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301
Charge-transfer salts with mononuclear and dinuclear ylide gold(I) complexes: X-ray structure of $[Au(CH_2PPh_3)_2](TCNQ)$ (TCNQ = 7,7′,8,8′-tetracyanoquinodimethane) (E. Cerrada, M.C. Gimeno, A. Laguna, M. Laguna, V. Orera and P.G. Jones), 203
Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyldiphenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of $[Cr(CO)_4]_2$TPTA and $[Cr(CO)_5]_4$TPTA (H. Meider, D. Matković-Čalogović and P. Planinić), 265
Cluster chemistry XCVIII. Formation of an Ru_5–dicarbon complex: X-ray structures of $Ru_5(\mu_5$-$C_2)(\mu$-$PPh_2)_2(\mu$-$Cl)_2(CO)_{11}$ and $Ru_5(\mu$-$H)(\mu_5$-$CC(PPh_2))(\mu$-$Cl)(\mu$-$PPh_2)(CO)_{12}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 191

Crystal and molecular structures of 1,2;3,4-Di-μ-*o*-methyl/*o*-chlorobenzoato-*OO'*-1,3-bis(o-methyl/o-chlorobenzoato-*O*)-1,2,4;2,3,4-di-μ_3-oxotetrakis[di-*n*-butyltin(IV)] and molecular structure of 1,2;3,4-Di-μ-*o*-hydroxybenzoato-*OO'*-1,3-bis(o-hydroxybenzoato-*O*)-1,2,4;2,3,4-di-μ_3-oxotetrakis[di-*n*-butyltin(IV)] compounds (S.P. Narula, S. Kaur, R. Shankar, S.K. Bharadwaj and R.K. Chadha), 181

Cyclometallated complexes of Pd(II) and Pt(II) with 2-phenylimidazoline (C. Navarro-Ranninger, F. Zamora, I. López-Solera, A. Monge and J.R. Masaguer), 149

Dicarbonylmolybdenum and tungsten complexes of α,β-unsaturated esters. Preparation, properties and application in the complexation of α,β-unsaturated ketones (Th. Schmidt), 139

Novel zirconium complexes derived from C_2-symmetric diamide ligands; the X-ray crystal structure of $[Zr(\eta^1\text{-}CH_2Ph)(\eta^2\text{-}CH_2Ph)\{(C_6H_3)_2\text{-}2,2'\text{-}(NCH_2C_6H_4Ph\text{-}4)_2\text{-}6,6'\text{-}Me_2\}]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343

Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $PCpRe(CO)_3$ and of the derivative $PCpReNO(CH_3)PPh_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273

Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[CpMo(CO)_3H]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(C_5H_5)Mo(CO)(\mu\text{-}SMe)_3Mo(CO)_2(\mu\text{-}SMe)Mo(CO)_2(C_5H_5)]$ (S. Poder-Guillou, P. Schollhammer, F.Y. Pétillon, J. Talarmin, S.E. Girdwood and K.W. Muir), 321

Schwermetall-π-Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293

Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(C_5H_5)_2Yb(\mu\text{-}SCH_2CH_2CH_2CH_3)]_2$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25

Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[HOs_3(CO)_{10}(O_2CC_5H_4FeC_5H_5)]$ and $[\{Os_3H(CO)_{10}\}_2(O_2CC_5H_4FeC_5H_4CO_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77

Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and −2,2'-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

Synthesis, spectroscopy and structure of phenylated cyclopentasilanes (U. Pöschl, H. Siegl and K. Hassler), 93

The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129

Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D. Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253

C–Si bond cleavage

A convenient preparation of pentafluorophenyl(fluoro)silanes: reactivity of pentafluorophenyltrifluorosilane (H.J. Frohn, M. Giesen, A. Klose, A. Lewin and V.V. Bardin), 155

Cyclometallation

Cyclometallated complexes of Pd(II) and Pt(II) with 2-phenylimidazoline (C. Navarro-Ranninger, F. Zamora, I. López-Solera, A. Monge and J.R. Masaguer), 149

Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortigueira), 165

Cyclopentadienyl

Ein ionischer Ti^{IV}–Fulvalenkomplex: $[\{CpTi(H_2O)\}_2(C_{10}H_8, O)]^{2+}(SO_3CF_3^-)_2 \cdot 1/2THF$ (U. Thewalt und T. Wöhrle), 331

Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $PCpRe(CO)_3$ and of the derivative $PCpReNO(CH_3)PPh_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273

Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121

Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[CpMo(CO)_3H]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(C_5H_5)Mo(CO)(\mu\text{-}SMe)_3Mo(CO)_2(\mu\text{-}SMe)Mo(CO)_2(C_5H_5)]$ (S. Poder-Guillou, P. Schollhammer, F.Y. Pétillon, J. Talarmin, S.E. Girdwood and K.W. Muir), 321

Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(C_5H_5)_2Yb(\mu\text{-}SCH_2CH_2CH_2CH_3)]_2$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25

Thermolysis of $(MeCp)_2Nb(\eta^2\text{-}CO_2)R$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49

Decarbonylation

Thermolysis of $(MeCp)_2Nb(\eta^2\text{-}CO_2)R$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49

Dialkylgallanes

Association and exchange in dialkylgallanes in toluene solution (D.K. Russell, A.S. Grady, R.E. Linney, Z. Mahmood and R.D. Markwell), 327

Diamide ligands

Novel zirconium complexes derived from C_2-symmetric diamide ligands; the X-ray crystal structure of $[Zr(\eta^1\text{-}CH_2Ph)(\eta^2\text{-}CH_2Ph)\{(C_6H_3)_2\text{-}2,2'\text{-}(NCH_2C_6H_4Ph\text{-}4)_2\text{-}6,6'\text{-}Me_2\}]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343

Dicarbon complex

Cluster chemistry XCVIII. Formation of an Ru_5–dicarbon complex: X-ray structures of $Ru_5(\mu_5\text{-}C_2)(\mu\text{-}PPh_2)_2(\mu\text{-}Cl)_2(CO)_{11}$ and $Ru_5(\mu\text{-}H)(\mu_3\text{-}CC(PPh_2))(\mu\text{-}Cl)(\mu\text{-}PPh_2)(CO)_{12}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 191

Diimine

Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortigueira), 165

Diruthenium complexes

Comparative chemistry of $\mu\text{-}\eta^1:\eta^2$-allenyl and $\mu\text{-}\eta^1:\eta^2$-acetylide complexes: η^5-cyclopentadienyl complexes via alkyne-allenyl coupling in the reactions of PhC≡CPh with $[Ru_2(CO)_6\{\mu\text{-}\eta^1:\eta^2_{\beta,\gamma}\text{-}C(Ph)=C=CH_2\}(\mu\text{-}PPh_2)]$ and $[Ru_2(CO)_6\{\mu\text{-}\eta^1:\eta^2\text{-}C=CPh\}(\mu\text{-}PPh_2)]$. X-Ray structures of $[\{Ru_2(CO)_4\{\mu\text{-}\eta^2(C,O),\eta^5\text{-}C_5MePh_2(C_6H_4)O\}(\mu\text{-}PPh_2)\}_2]$ and $[Ru_2(CO)_5\{\mu\text{-}\eta^2(C,O),\eta^5\text{-}C_5HPh_2(C_6H_4)O\}(\mu\text{-}PPh_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229

Dysprosium

Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(C_5H_5)_2Yb(\mu\text{-}$

$SCH_2CH_2CH_2CH_3)]_2$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25

Electrochemistry
Redox behaviour of ferrocene derivatives. VIII. 1,1'-Bis(diphenylphosphino)ferrocenes (P. Zanello, G. Opromolla, G. Giorgi, G. Sasso and A. Togni), 61
The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129

Electronic spectra
Photoredox decomposition of the adduct of methyltrioxorhenium(VII) and hexacyanoferrate(II) induced by metal-to-metal charge transfer excitation (H. Kunkely and A. Vogler), 175

Electron Spin resonance
A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: $[(C_5HMe_4)_2Ti(\eta^1\text{-}C{\equiv}CC{\equiv}CSiMe_3)_2]^-$-$[Li(THF)_2]^+$ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109

Enantioselective catalysis
Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177

Exchange
Association and exchange in dialkylgallanes in toluene solution (D.K. Russell, A.S. Grady, R.E. Linney, Z. Mahmood and R.D. Markwell), 327

Ferricinium
Synthesis and characterization of new perfluoroalkylated side-chain ferrocenes and ferricinium salts (C. Guillon and P. Vierling), 211

Ferrocene
The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129

Ferrocenes
Redox behaviour of ferrocene derivatives. VIII. 1,1'-Bis(diphenylphosphino)ferrocenes (P. Zanello, G. Opromolla, G. Giorgi, G. Sasso and A. Togni), 61
Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[HOs_3(CO)_{10}(O_2CC_5H_4FeC_5H_5)]$ and $[\{Os_3H(CO)_{10}\}_2(O_2CC_5H_4FeC_5H_4CO_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77

Ferrocenyl alcohols
The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129

Fluoride binding constant
Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85

Fluorine Silicon
Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85

Fluorophoshine-complexes
Übergangsmetall-Trifluormethyl-Komplexe: Erste Trifluormethyl-Komplexe von Gold mit Fluorphosphan-Liganden (A. Gräfe und T. Kruck), 31

Fulvalene bridge
Ein ionischer Ti^{IV}–Fulvalenkomplex: $[[CpTi(H_2O)]_2(C_{10}H_8, O)]^{2+}(SO_3CF_3^-)_2 \cdot 1/2THF$ (U. Thewalt und T. Wöhrle), 331

Gallium
Association and exchange in dialkylgallanes in toluene solution (D.K. Russell, A.S. Grady, R.E. Linney, Z. Mahmood and R.D. Markwell), 327

Gold
Charge-transfer salts with mononuclear and dinuclear ylide gold(I) complexes: X-ray structure of $[Au(CH_2PPh_3)_2](TCNQ)$ (TCNQ = 7,7',8,8'-tetracyanoquinodimethane) (E. Cerrada, M.C. Gimeno, A. Laguna, M. Laguna, V. Orera and P.G. Jones), 203
Übergangsmetall-Trifluormethyl-Komplexe: Erste Trifluormethyl-Komplexe von Gold mit Fluorphosphan-Liganden (A. Gräfe und T. Kruck), 31

Hafnium
Unsymmetrical *ansa*-metallocenes of zirconium and hafnium (W.A. Herrmann, M.J.A. Morawietz and T. Priermeier), 351

Hydroformylation, 3-vinylpyrrole, 2-(3-pyrrolyl)propanal
High α-regioselectivity in the rhodium-catalyzed hydroformylation of vinylpyrroles (R. Settambolo, A. Caiazzo and R. Lazzaroni), 337

Hydroxyacetophenone
Ruthenium-catalysed *ortho* alkylation of hydroxyacetophenones; the functionalisation of ring C aromatic diterpenoids (P.W.R. Harris and P.D. Woodgate), 339

Hypervalent
Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin–carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259

Insertion
Thermolysis of $(MeCp)_2Nb(\eta^2\text{-}CO_2)R$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49

Iridium
Direct high yield synthesis of $[Ir(OH)(CO)(PPh_3)_2]$ from Vaska's compound by phase-transfer catalysis (S.A. Al-Jibori), 119

Iron
A chalcogen-stabilized cluster containing a *s-trans*-1,3-diene ligand: synthesis and structural characterization of $\{(CO)_6Fe_2Se_2\}_2(\mu\text{-}s\text{-}trans\text{-}C_4H_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307
Photoredox decomposition of the adduct of methyltrioxorhenium(VII) and hexacyanoferrate(II) induced by metal-to-metal charge transfer excitation (H. Kunkely and A. Vogler), 175
Propargyl alcohol derivatives of $Fe_2(CO)_6(\mu\text{-}EE')$ (E,E' = S, Se or Te): synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu\text{-}EC(H){=}C(CH_2OH)E'\}$ (E = Se or Te; E' = S, Se or Te); Crystal structure of $(CO)_5(PPh_3)Fe_2\{\mu\text{-}SeC(CH_2OH){=}C(H)Se\}$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313
Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121
Reaktionen von Nitrosylkomplexen. XII. Umsetzung von $[\{C_5Me_5\text{-}Fe(\mu\text{-}NO)\}_2]$ mit $[C_5Me_5Co(C_2H_4)_2]$: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnoukh), 113
Redox behaviour of ferrocene derivatives. VIII. 1,1'-Bis(diphenylphosphino)ferrocenes (P. Zanello, G. Opromolla, G. Giorgi, G. Sasso and A. Togni), 61
Synthesis and characterization of new perfluoroalkylated side-chain ferrocenes and ferricinium salts (C. Guillon and P. Vierling), 211
Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[HOs_3(CO)_{10}(O_2CC_5H_4FeC_5H_5)]$ and $[\{Os_3H(CO)_{10}\}_2(O_2CC_5H_4FeC_5H_4CO_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77
The redox behaviour of ferrocenyl alcohols and ferrocenediyl-diols: crystal and molecular structure of ferrocenyl(2-furyl)phenylmethanol (G. Ferguson, C. Glidewell, G. Opromolla, C.M. Zakaria and P. Zanello), 129

IR spectroscopy
Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301
Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $PCpRe(CO)_3$ and of the derivative $PCpReNO(CH_3)PPh_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273

Kinetic data
New rhodium(I)-π-chelate complexes with coordinated amidine bases (dbu, dbn) and their catalytic properties to polymerize phenylacetylene (J. Schniedermeier and H.-J. Haupt), 41

Kinetic isotope effects
α-Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Lanthanides
The first structurally authenticated divalent lanthanide siloxide derivative, $[Yb(OSiMe_2{}^tBu)(\eta^2\text{-dme})(\mu\text{-}OSiMe_2{}^tBu)]_2$ (D.J. Duncalf, P.B. Hitchcock and G.A. Lawless), 347

Lanthanoid
Complexes of lanthanoids with neutral π donor ligands (G.B. Deacon and Q. Shen), 1

Ligand exchange
Dicarbonylmolybdenum and tungsten complexes of α,β-unsaturated esters. Preparation, properties and application in the complexation of α,β-unsaturated ketones (Th. Schmidt), 139

Lithium
A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: $[(C_5HMe_4)_2Ti(\eta^1\text{-}C{\equiv}CC{\equiv}CSiMe_3)_2]^-\text{-}[Li(THF)_2]^+$ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109
α-Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Macrocycle
Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyldiphenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of $[Cr(CO)_4]_2TPTA$ and $[Cr(CO)_5]_4TPTA$ (H. Meider, D. Matković-Čalogović and P. Planinić), 265

Magnesium
α-Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Manganese
Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortigueira), 165

Mass spectrometry
Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301
Redox behaviour of ferrocene derivatives. VIII. 1,1'-Bis(diphenylphosphino)ferrocenes (P. Zanello, G. Opromolla, G. Giorgi, G. Sasso and A. Togni), 61

Metal amides
Unsymmetrical *ansa*-metallocenes of zirconium and hafnium (W.A. Herrmann, M.J.A. Morawietz and T. Priermeier), 351

Methyllithium
α-Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Methylmagnesium iodide
α-Deuterium kinetic isotope effects in reactions of methyllithium: is better aggregation the cause of lower reactivity? (T. Holm), 37

Molecular tweezer
A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: $[(C_5HMe_4)_2Ti(\eta^1\text{-}C{\equiv}CC{\equiv}CSiMe_3)_2]^-\text{-}[Li(THF)_2]^+$ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109

Molybdenum
Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301
Dicarbonylmolybdenum and tungsten complexes of α,β-unsaturated esters. Preparation, properties and application in the complexation of α,β-unsaturated ketones (Th. Schmidt), 139
Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[CpMo(CO)_3H]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(C_5H_5)Mo(CO)(\mu\text{-}SMe)_3Mo(CO)_2(\mu\text{-}SMe)Mo(CO)_2(C_5H_5)]$ (S. Poder-Guillou, P. Schollhammer, F.Y. Pétillon, J. Talarmin, S.E. Girdwood and K.W. Muir), 321

Mössbauer spectra
Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and $-2,2'$-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

Mössbauer spectroscopy
Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D. Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253

Niobium
Bis-η-cyclopentadienyl complexes of niobium and tungsten with silyl and stannyl ligands (M.L.H. Green and A.K. Hughes), 221
First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357
Thermolysis of $(MeCp)_2Nb(\eta^2\text{-}CO_2)R$ complexes: decarbonylation vs. insertion (P.-F. Fu, M.A. Khan and K.M. Nicholas), 49

Nitrogen bases
Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and $-2,2'$-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

Nitrosyl complexes
Reaktionen von Nitrosylkomplexen. XII. Umsetzung von $[(C_5Me_5\text{-}Fe(\mu\text{-}NO))_2]$ mit $[C_5Me_5Co(C_2H_4)_2]$: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnoukh), 113

NMR
Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2'-bipyrimidine and further NMR studies of diorganotin-pyrazine and $-2,2'$-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67

NMR spectroscopy
Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301
Dicarbonylmolybdenum and tungsten complexes of α,β-unsaturated esters. Preparation, properties and application in the complexation of α,β-unsaturated ketones (Th. Schmidt), 139
Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $PCpRe(CO)_3$ and of the derivative $PCpReNO(CH_3)PPh_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273

Nuclear magnetic resonance
Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D. Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253
Nucleophilic arylation of electrophiles
A convenient preparation of pentafluorophenyl(fluoro)silanes: reactivity of pentafluorophenyltrifluorosilane (H.J. Frohn, M. Giesen, A. Klose, A. Lewin and V.V. Bardin), 155
Nucleophilic substitution
Synthesis and characterization of tricarbonyl[η^6-(phenylethynyl)-arene]chromium complexes: crystal structure of tricarbonyl[η^6-(3-phenylethynyl)anisole]chromium (F.-E. Hong, S.-C. Lo, M.-W. Liou, Y.-T. Chang and C.-C. Lin), 101

Olefin polymerization
Novel zirconium complexes derived from C_2-symmetric diamide ligands; the X-ray crystal structure of $[Zr(\eta^1\text{-}CH_2Ph)(\eta^2\text{-}CH_2Ph)\{(C_6H_3)_2\text{-}2,2'\text{-}(NCH_2C_6H_4Ph\text{-}4)_2\text{-}6,6'\text{-}Me_2\}]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343
Olefins
Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177
Optically active ligands
Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $PCpRe(CO)_3$ and of the derivative $PCpReNO(CH_3)PPh_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273
Organolanthoid chemistry
Complexes of lanthanoids with neutral π donor ligands (G.B. Deacon and Q. Shen), 1
Organotin
Synthesis, spectroscopic (Mössbauer, IR and NMR) and X-ray structural studies of diorganotin complexes of 2,2′-bipyrimidine and further NMR studies of diorganotin-pyrazine and −2,2′-azopyridine complexes (F. Caruso, M. Giomini, A.M. Giuliani and E. Rivarola), 67
Ortho alkylation
Ruthenium-catalysed *ortho* alkylation of hydroxyacetophenones; the functionalisation of ring C aromatic diterpenoids (P.W.R. Harris and P.D. Woodgate), 339
Osmium
Synthesis, characterization, electrochemical properties and crystal and molecular structures of $[HOs_3(CO)_{10}(O_2CC_5H_4FeC_5H_5)]$ and $[\{Os_3H(CO)_{10}\}_2(O_2CC_5H_4FeC_5H_4CO_2)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), 77
2-Oxoglutarates
Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177

Palladium
Cyclometallated complexes of Pd(II) and Pt(II) with 2-phenylimidazoline (C. Navarro-Ranninger, F. Zamora, I. López-Solera, A. Monge and J.R. Masaguer), 149
Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortigueira), 165
Palladium complexes
Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177
π donor ligands
Complexes of lanthanoids with neutral π donor ligands (G.B. Deacon and Q. Shen), 1
Pentacoordinate
Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin–carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259
Pentacoordinate siliconate
Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85
Pentafluorophenylhalogenosilanes
A convenient preparation of pentafluorophenyl(fluoro)silanes: reactivity of pentafluorophenyltrifluorosilane (H.J. Frohn, M. Giesen, A. Klose, A. Lewin and V.V. Bardin), 155
Pentamethylcyclopentadienyl complexes
Reaktionen von Nitrosylkomplexen. XII. Umsetzung von $[\{C_5Me_5\text{-}Fe(\mu\text{-}NO)\}_2]$ mit $[C_5Me_5Co(C_2H_4)_2]$: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnoukh), 113
Perfluoroalkylated ferrocene
Synthesis and characterization of new perfluoroalkylated side-chain ferrocenes and ferricinium salts (C. Guillon and P. Vierling), 211
Phase transfer catalysis
Direct high yield synthesis of $[Ir(OH)(CO)(PPh_3)_2]$ from Vaska's compound by phase-transfer catalysis (S.A. Al-Jibori), 119
Phenylated cyclopentasilanes
Synthesis, spectroscopy and structure of phenylated cyclopentasilanes (U. Pöschl, H. Siegl and K. Hassler), 93
Phenylethynylarene
Synthesis and characterization of tricarbonyl[η^6-(phenylethynyl)-arene]chromium complexes: crystal structure of tricarbonyl[η^6-(3-phenylethynyl)anisole]chromium (F.-E. Hong, S.-C. Lo, M.-W. Liou, Y.-T. Chang and C.-C. Lin), 101
Phosphinoalkyne
Reactions of cationic iron P-coordinated (diphenylphosphino)alkynes with dicobalt octacarbonyl (E. Louattani, J. Suades, A. Alvarez-Larena, J.F. Piniella and G. Germain), 121
Photochemistry
Photoredox decomposition of the adduct of methyltrioxorhenium(VII) and hexacyanoferrate(II) induced by metal-to-metal charge transfer excitation (H. Kunkely and A. Vogler), 175
Platinum
Cyclometallated complexes of Pd(II) and Pt(II) with 2-phenylimidazoline (C. Navarro-Ranninger, F. Zamora, I. López-Solera, A. Monge and J.R. Masaguer), 149
Polymerization of phenylacetylene
New rhodium(I)-π-chelate complexes with coordinated amidine bases (dbu, dbn) and their catalytic properties to polymerize phenylacetylene (J. Schniedermeier and H.-J. Haupt), 41
Propargyl alcohol
Propargyl alcohol derivatives of $Fe_2(CO)_6(\mu\text{-}EE')$ (E,E′ = S, Se or Te): synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu\text{-}EC(H)=C(CH_2OH)E'\}$ (E = Se or Te; E′ = S, Se or Te); Crystal structure of $(CO)_5(PPh_3)Fe_2\{\mu\text{-}SeC(CH_2OH)=C(H)Se\}$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313

X-Ray structure
Ein ionischer Ti^{IV}–Fulvalenkomplex: $[[CpTi(H_2O)]_2(C_{10}H_8, O)]^{2+}(SO_3CF_3^-)_2 \cdot 1/2THF$ (U. Thewalt und T. Wöhrle), 331
Rearrangement
First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357
Rhenium
Optically active transition-metal complexes. II. Rhenium complexes with the optically active cyclopentadienyl ligand PCp: X-ray structures of the *exo* and *endo* isomers of $PCpRe(CO)_3$ and of the

derivative $PCpReNO(CH_3)PPh_3$ (W.H. Bosch, U. Englert, B. Pfister, R. Stauber and A. Salzer), 273

Photoredox decomposition of the adduct of methyltrioxorhenium(VII) and hexacyanoferrate(II) induced by metal-to-metal charge transfer excitation (H. Kunkely and A. Vogler), 175

Rh(I) complexes

New rhodium(I)-π-chelate complexes with coordinated amidine bases (dbu, dbn) and their catalytic properties to polymerize phenylacetylene (J. Schniedermeier and H.-J. Haupt), 41

Rhodium

High α-regioselectivity in the rhodium-catalyzed hydroformylation of vinylpyrroles (R. Settambolo, A. Caiazzo and R. Lazzaroni), 337

Ruthenium

Cluster chemistry XCVIII. Formation of an Ru_3–dicarbon complex: X-ray structures of $Ru_5(\mu_5\text{-}C_2)(\mu\text{-}PPh_2)_2(\mu\text{-}Cl)_2(CO)_{11}$ and $Ru_5(\mu\text{-}H)(\mu_5\text{-}CC(PPh_2))(\mu\text{-}Cl)(\mu\text{-}PPh_2)(CO)_{12}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), 191

Comparative chemistry of $\mu\text{-}\eta^1:\eta^2$-allenyl and $\mu\text{-}\eta^1:\eta^2$-acetylide complexes: η^5-cyclopentadienyl complexes via alkyne-allenyl coupling in the reactions of PhC≡CPh with $[Ru_2(CO)_6(\mu\text{-}\eta^1:\eta^2_{3,4}\text{-}C(Ph)=C=CH_2)(\mu\text{-}PPh_2)]$ and $[Ru_2(CO)_6(\mu\text{-}\eta^1:\eta^2\text{-}C\equiv CPh)(\mu\text{-}PPh_2)]$. X-Ray structures of $[\{Ru_2(CO)_4(\mu\text{-}\eta^2(C,O),\eta^5\text{-}C_5MePh_2(C_6H_4)(O))(\mu\text{-}PPh_2)\}_2]$ and $[Ru_2(CO)_5\{\mu\text{-}\eta^2(C,O),\eta^5\text{-}C_5HPh_2(C_6H_4)(O)\}(\mu\text{-}PPh_2)]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229

Ruthenium-catalysed *ortho* alkylation of hydroxyacetophenones; the functionalisation of ring C aromatic diterpenoids (P.W.R. Harris and P.D. Woodgate), 339

Schiff bases

Synthesis and characterization of cyclometallated complexes of palladium(II) and manganese(I) with bidentate Schiff bases (J.M. Vila, M. Gayoso, T. Pereira, M. López Torres, J.J. Fernández, A. Fernández and J.M. Ortigueira), 165

Selenium

A chalcogen-stabilized cluster containing a *s-trans*-1,3-diene ligand: synthesis and structural characterization of $[(CO)_6Fe_2Se_2]_2(\mu\text{-}s\text{-}trans\text{-}C_4H_2)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana and B. Verghese), 307

Propargyl alcohol derivatives of $Fe_2(CO)_6(\mu\text{-}EE')$ (E,E' = S, Se or Te): synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu\text{-}EC(H)=C(CH_2OH)E'\}$ (E = Se or Te; E' = S, Se or Te); Crystal structure of $(CO)_5(PPh_3)Fe_2\{\mu\text{-}SeC(CH_2OH)=C(H)Se\}$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313

Silane

Anion complexation by bidentate Lewis acidic hosts, *ortho*-bis(fluorosilyl)benzenes (K. Tamao, T. Hayashi and Y. Ito), 85

Silicon

Bis-η-cyclopentadienyl complexes of niobium and tungsten with silyl and stannyl ligands (M.L.H. Green and A.K. Hughes), 221

The first structurally authenticated divalent lanthanide siloxide derivative, $[Yb(OSiMe_2{}^tBu)(\eta^2\text{-}dme)(\mu\text{-}OSiMe_2{}^tBu)]_2$ (D.J. Duncalf, P.B. Hitchcock and G.A. Lawless), 347

Siloxide

The first structurally authenticated divalent lanthanide siloxide derivative, $[Yb(OSiMe_2{}^tBu)(\eta^2\text{-}dme)(\mu\text{-}OSiMe_2{}^tBu)]_2$ (D.J. Duncalf, P.B. Hitchcock and G.A. Lawless), 347

Si–N Bond cleavage

First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357

^{29}Si NMR spectroscopy

Synthesis, spectroscopy and structure of phenylated cyclopentasilanes (U. Pöschl, H. Siegl and K. Hassler), 93

Sulfur

Propargyl alcohol derivatives of $Fe_2(CO)_6(\mu\text{-}EE')$ (E,E' = S, Se or Te): synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu\text{-}EC(H)=C(CH_2OH)E'\}$ (E = Se or Te; E' = S, Se or Te); Crystal structure of $(CO)_5(PPh_3)Fe_2\{\mu\text{-}SeC(CH_2OH)=C(H)Se\}$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313

Tantalum

First amido-functionalized niobium and tantalum complexes of the *ansa*-structural type: synthesis and photochemical Si-N bond cleavage (W.A. Herrmann and W. Baratta), 357

Tellurium

Propargyl alcohol derivatives of $Fe_2(CO)_6(\mu\text{-}EE')$ (E,E' = S, Se or Te): synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu\text{-}EC(H)=C(CH_2OH)E'\}$ (E = Se or Te; E' = S, Se or Te); Crystal structure of $(CO)_5(PPh_3)Fe_2\{\mu\text{-}SeC(CH_2OH)=C(H)Se\}$ (P. Mathur, Md.M. Hossain, S. Umbarkar, C.V.V. Satyanarayana and V.G. Puranik), 313

Tertiary stibines

Tetrachloropyridylstibines (P. Sharma and N.K. Jha), 19

Tetraaza ligand

Chromium carbonyl complexes containing 1,4,8,11-tetrakis(methyldiphenylphosphino)-1,4,8,11-tetraazacyclotetradecane (TPTA): the synthesis and characterization of $[Cr(CO)_4]_2TPTA$ and $[Cr(CO)_5]_4TPTA$ (H. Meider, D. Matković-Čalogović and P. Planinić), 265

Tetrachloroaluminates

Schwermetall-π-Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293

Tetrachlorogallates

Schwermetall-π-Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293

Tetrachloropyridyl stibines

Tetrachloropyridylstibines (P. Sharma and N.K. Jha), 19

Thallium–arene π-bonding

Schwermetall-π-Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293

Thallium complexes

Schwermetall-π-Komplexe. XI. Synthese, Struktur und thermisches Verhalten von Bis(1,2,4-trimethylbenzol)thallium(I)-tetrachloroaluminat bzw. -gallat — Zum Einfluß des Gegenions auf Struktur und Stabilität eines kationischen Bis(aren)hauptgruppenmetallkomplexes (W. Frank, G. Korrell und G.J. Reiß), 293

Thermolysis

Titanocene-bis(trimethylsilyl)acetylene complexes: effects of methyl substituents at the cyclopentadienyl ligands on the structure of thermolytic products (V. Varga, K. Mach, M. Polášek, P. Sedmera, J. Hiller, U. Thewalt and S.I. Troyanov), 241

Thiazole

Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D.

Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253

Thiolate
Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(C_5H_5)_2Yb(\mu\text{-}SCH_2CH_2CH_2CH_3)]_2$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25

Tin
Bis-η-cyclopentadienyl complexes of niobium and tungsten with silyl and stannyl ligands (M.L.H. Green and A.K. Hughes), 221
Crystal and molecular structures of 1,2;3,4-Di-μ-*o*-methyl/*o*-chlorobenzoato-*OO'*-1,3-bis(o-methyl/o-chlorobenzoato-*O*)-1,2,4;2,3,4-di-μ_3-oxotetrakis[di-*n*-butyltin(IV)] and molecular structure of 1,2;3,4-Di-μ-*o*-hydroxybenzoato-*OO'*-1,3-bis(o-hydroxybenzoato-*O*)-1,2,4;2,3,4-di-μ_3-oxotetrakis[di-*n*-butyltin(IV)] compounds (S.P. Narula, S. Kaur, R. Shankar, S.K. Bharadwaj and R.K. Chadha), 181
Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin–carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259
Thiazole complexes of diorganotin(IV) dihalides: the crystal structure of dichlorodiphenylbis(thiazole)tin(IV) (P. Alvarez Boo, M.D. Couce, E. Freijanes, J.S. Casas, A. Castiñeiras, A. Sánchez González, J. Sordo and U. Russo), 253

Tin ate complex
Synthesis, stability, and reactions of 10-Sn-5 tin ate complexes bearing four tin–carbon bonds (Y. Yamamoto, A. Sakaguchi, N. Ohashi and K.-Y. Akiba), 259

Titanium
Ein ionischer Ti^{IV}–Fulvalenkomplex: $[[CpTi(H_2O)]_2(C_{10}H_8, O)]^{2+}(SO_3CF_3^-)_2 \cdot 1/2THF$ (U. Thewalt und T. Wöhrle), 331
Titanocene-bis(trimethylsilyl)acetylene complexes: effects of methyl substituents at the cyclopentadienyl ligands on the structure of thermolytic products (V. Varga, K. Mach, M. Polášek, P. Sedmera, J. Hiller, U. Thewalt and S.I. Troyanov), 241

Titanium (III)
A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: $[(C_5HMe_4)_2Ti(\eta^1\text{-}C{\equiv}CC{\equiv}CSiMe_3)_2]^-[Li(THF)_2]^+$ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109

Titanocene
Titanocene-bis(trimethylsilyl)acetylene complexes: effects of methyl substituents at the cyclopentadienyl ligands on the structure of thermolytic products (V. Varga, K. Mach, M. Polášek, P. Sedmera, J. Hiller, U. Thewalt and S.I. Troyanov), 241

Triazolylborate
Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301

Tricarbonyl
Carbonyl–molybdenum complexes of the hydrotris(3,5-dimethyl-1,2,4-triazol-1-yl)borate ligand (I.T. Macleod, E.R.T. Tiekink and C.G. Young), 301

Trifluoromethane sulphonate
Ein ionischer Ti^{IV}–Fulvalenkomplex: $[[CpTi(H_2O)]_2(C_{10}H_8, O)]^{2+}(SO_3CF_3^-)_2 \cdot 1/2THF$ (U. Thewalt und T. Wöhrle), 331

Trifluoromethyl-complexes
Übergangsmetall-Trifluormethyl-Komplexe: Erste Trifluormethyl-Komplexe von Gold mit Fluorphosphan-Liganden (A. Gräfe und T. Kruck), 31

Trinuclear complex
Reactions of dinuclear and polynuclear complexes. XV. Reinvestigation of the reaction between $[CpMo(CO)_3H]$, allyl chloride and dimethyl disulfide: Crystal structure of $[(C_5H_5)Mo(CO)(\mu\text{-}SMe)_3Mo(CO)_2(\mu\text{-}SMe)Mo(CO)_2(C_5H_5)]$ (S. Poder-Guillou, P. Schollhammer, F.Y. Pétillon, J. Talarmin, S.E. Girdwood and K.W. Muir), 321

Triple carbonylation
Catalytic triple carbonylation of olefins. Enantioselective synthesis of 2-oxoglutarates (M. Sperrle and G. Consiglio), 177

Tungsten
Bis-η-cyclopentadienyl complexes of niobium and tungsten with silyl and stannyl ligands (M.L.H. Green and A.K. Hughes), 221
Dicarbonylmolybdenum and tungsten complexes of α,β-unsaturated esters. Preparation, properties and application in the complexation of α,β-unsaturated ketones (Th. Schmidt), 139

α,β-Unsaturated esters
Dicarbonylmolybdenum and tungsten complexes of α,β-unsaturated esters. Preparation, properties and application in the complexation of α,β-unsaturated ketones (Th. Schmidt), 139

Vaska's compound
Direct high yield synthesis of $[Ir(OH)(CO)(PPh_3)_2]$ from Vaska's compound by phase-transfer catalysis (S.A. Al-Jibori), 119

X-ray crystallography
Reaktionen von Nitrosylkomplexen. XII. Umsetzung von $[\{C_5Me_5Fe(\mu\text{-}NO)\}_2]$ mit $[C_5Me_5Co(C_2H_4)_2]$: NO transfer vs. Cycloaddition (J. Müller, I. Sonn und T. Akhnoukh), 113

X-ray diffraction
A titanium(III) tweezer complex with an embedded alkali metal ion between diynyl ligands: $[(C_5HMe_4)_2Ti(\eta^1\text{-}C{\equiv}CC{\equiv}CSiMe_3)_2]^-[Li(THF)_2]^+$ (V. Varga, K. Mach, J. Hiller and U. Thewalt), 109

X-ray structures
Comparative chemistry of $\mu\text{-}\eta^1:\eta^2$-allenyl and $\mu\text{-}\eta^1:\eta^2$-acetylide complexes: η^5-cyclopentadienyl complexes via alkyne-allenyl coupling in the reactions of PhC≡CPh with $[Ru_2(CO)_6(\mu\text{-}\eta^1:\eta^2_{\beta,\gamma}\text{-}C(Ph)=C=CH_2)(\mu\text{-}PPh_2)]$ and $[Ru_2(CO)_6(\mu\text{-}\eta^1:\eta^2\text{-}C{\equiv}CPh)(\mu\text{-}PPh_2)]$. X-Ray structures of $[\{Ru_2(CO)_4(\mu\text{-}\eta^2(C,O),\eta^5\text{-}C_5MePh_2(C_6H_4)O)(\mu\text{-}PPh_2)\}_2]$ and $[Ru_2(CO)_5\{\mu\text{-}\eta^2(C,O),\eta^5\text{-}C_5HPh_2(C_6H_4)O)(\mu\text{-}PPh_2)\}]$ (P. Blenkiron, S.M. Breckenridge, N.J. Taylor, A.J. Carty, M.A. Pellinghelli, A. Tiripicchio and E. Sappa), 229

Ylide
Charge-transfer salts with mononuclear and dinuclear ylide gold(I) complexes: X-ray structure of $[Au(CH_2PPh_3)_2](TCNQ)$ (TCNQ = 7,7',8,8'-tetracyanoquinodimethane) (E. Cerrada, M.C. Gimeno, A. Laguna, M. Laguna, V. Orera and P.G. Jones), 203

Ytterbium
Synthesis and thermal stability of binuclear biscyclopentadienyl lanthanide thiolates, X-ray crystal structure of $[(C_5H_5)_2Yb(\mu\text{-}SCH_2CH_2CH_2CH_3)]_2$ (Z.-Z. Wu, Z.-E. Huang, R.-F. Cai, X.-G. Zhou, Z. Xu, X.-Z. You and X.-Y. Huang), 25
The first structurally authenticated divalent lanthanide siloxide derivative, $[Yb(OSiMe_2{}^tBu)(\eta^2\text{-dme})(\mu\text{-}OSiMe_2{}^tBu)]_2$ (D.J. Duncalf, P.B. Hitchcock and G.A. Lawless), 347

Zirconium
Novel zirconium complexes derived from C_2-symmetric diamide ligands; the X-ray crystal structure of $[Zr(\eta^1\text{-}CH_2Ph)(\eta^2\text{-}CH_2Ph)\{(C_6H_3)_2\text{-}2,2'\text{-}(NCH_2C_6H_4Ph\text{-}4)_2\text{-}6,6'\text{-}Me_2\}]$ (F.G.N. Cloke, T.J. Geldbach, P.B. Hitchcock and J.B. Love), 343
Unsymmetrical *ansa*-metallocenes of zirconium and hafnium (W.A. Herrmann, M.J.A. Morawietz and T. Priermeier), 351